

Brunet-Derrida particle systems, free boundary problems and Wiener-Hopf equations

Rick Durrett and Daniel Remenik

Department of Mathematics and Center for Applied Mathematics
Cornell University, Ithaca, New York 14853

October 19, 2009

Abstract

We consider a branching-selection system in $\mathbb{R}$ with N particles which give birth independently at rate 1 and where after each birth the leftmost particle is erased, keeping the number of particles constant. We show that, as $N \rightarrow \infty$, the empirical measure process associated to the system converges in distribution to a deterministic measure-valued process whose densities solve a free boundary integro-differential equation. We also show that this equation has a unique traveling wave solution traveling at speed c or no such solution depending on whether $c \geq a$ or $c < a$, where a is the asymptotic speed of the branching random walk obtained by ignoring the removal of the leftmost particles in our process. The traveling wave solutions correspond to solutions of Wiener-Hopf equations.

1 Introduction and statement of the results

We will consider the following branching-selection particle system. At any time t we have N particles on the real line with positions $\eta_t^N(1) \geq \dots \geq \eta_t^N(N)$. Each one of the N particles gives birth at rate 1 to a new particle whose position is chosen, relative to the parent particle, using a given probability distribution ρ on $\mathbb{R}$. Whenever a new particle is born, we reorder the $N + 1$ particles and erase the leftmost one (so the number of particles is always kept equal to N). We will denote by $\mathcal{X}_N = \{(\xi(1), \dots, \xi(N)) \in \mathbb{R}^N : \xi(1) \geq \dots \geq \xi(N)\}$ the state space of our process.

We learned of this process through the work of Durrett and Mayberry (2008), who considered the special case in which ρ corresponds to a uniform random variable on $[-1, 1]$. However, our process is a member of a family of processes that first arose in work of Brunet and Derrida (1997), who studied discrete approximations of the Fisher-Kolmogorov PDE:

$$\frac{\partial h}{\partial t} = \frac{\partial^2 h}{\partial t^2} + h - h^3.$$

In their discretization they have N particles at each integer time k with integer positions $x_1(k), \dots, x_N(k)$. At each time k they choose, for each $i \in \{1, \dots, k\}$, two particles j_i and j'_i at random (and with replacement) from $\{1, \dots, N\}$ and set

$$x_i(k+1) = \max\{x_{j_i}(k) + \alpha_i, x_{j'_i}(k) + \alpha'_i\},$$

Both authors were partially supported by NSF grant DMS 0704996 from the probability program.

where $\alpha_i, \alpha'_i, 1 \leq i \leq N$, are independent $= 1$ with probability p and $= 0$ with probability $1 - p$ (and are resampled at each time step).

They conjectured that the propagation speed of the finite system converged to the minimal wave speed of the PDE at a slow rate, of order $(\log N)^{-2}$. Bérard and Gouéré (2008) considered a discrete time version of the process where at each time step each of the N particles are replaced by two new particles, and then only the N rightmost ones are kept. Under some assumptions on the distribution used to choose the locations of the new particles (a strong one being that new particles are always sent to the right of the parent particle) they proved that the system moves to the right with a deterministic asymptotic speed v_N which converges as $N \rightarrow \infty$ to some explicit v at the predicted rate.

Although we will say something about the behavior of the system for fixed N , our main interest in this paper is to study the behavior of the empirical distribution of the process as $N \rightarrow \infty$. Before proceeding with this, let us specify some assumptions. When a particle at x gives birth, the new particle is sent to a location $x + y$ with y being chosen from an absolutely continuous probability distribution $\rho(y)dy$. We will assume that ρ is symmetric and that $\int_0^\infty |x|\rho(x) dx < \infty$. The initial condition for our process will always be specified as follows: each particle starts at a location chosen independently from a probability measure $f_0(x)dx$, where $f_0(x) = 0$ for $x < 0$ and $f_0(x)$ is strictly positive and continuous for $x > 0$.

1.1 Convergence to the solution of a free boundary problem

Let

$$\nu_t^N = \frac{1}{N} \sum_{i=1}^N \delta_{\eta_t^N(i)}$$

be the empirical measure associated to η_t^N . Observe that the initial empirical measure $\nu_0^N(dx)$ converges in distribution to $f_0(x)dx$. We will show that, as $N \rightarrow \infty$, this empirical measure process converges to a deterministic measure-valued process whose densities are the solution of a certain free boundary problem.

Alternatively one could think of the following (weaker) version of the problem. It is not hard to see that the probability measure $\mathbb{E}(\nu_t^N(\cdot))$ on $\mathbb{R}$ is absolutely continuous. Letting $\bar{f}^N(t, x)$ be its density, then we want to study its limit as $N \rightarrow \infty$. We expect this limit $f(t, x)$ to correspond to the densities of the limiting measure-valued process mentioned above. Now observe that if ξ_t^N is a version of our process in which we do not erase the leftmost particle after births (i.e., ξ_t^N is a branching random walk), then we would expect that the density of the corresponding expected empirical measure converges to the solution $\hat{f}(t, x)$ of the following integro-differential equation:

$$(1.1) \quad \frac{\partial \hat{f}}{\partial t}(t, x) = \int_{-\infty}^{\infty} \hat{f}(t, y) \rho(x - y) dy.$$

This is indeed the case, as we will see in Proposition 2.1. Observe that the total mass of $\hat{f}(t, \cdot)$ grows exponentially (in fact $\int_{-\infty}^{\infty} \hat{f}(t, y) dy = e^t$).

By adding the selection step to our process we ensure that the limiting density always has mass 1, but otherwise the branching mechanism is still governed by the convolution

term appearing in (1.1). Thus we expect that if the limit $f(t, x) = \lim_{N \rightarrow \infty} \bar{f}^N(t, x)$ exists, then it has to satisfy the following: there exists a continuous non-decreasing function $\gamma: [0, \infty) \rightarrow \mathbb{R}$ with $\gamma(0) = 0$ such that $(f(t, x), \gamma(t))$ is the unique solution to the following free boundary problem (FB):

$$(FB1) \quad \frac{\partial f}{\partial t}(t, x) = \int_{-\infty}^{\infty} f(t, y) \rho(x - y) dy \quad \forall x > \gamma(t)$$

$$(FB2) \quad \int_{\gamma(t)}^{\infty} f(t, y) dy = 1$$

$$(FB3) \quad f(t, x) = 0 \quad \forall x \leq \gamma(t)$$

with initial condition $f(0, x) = f_0(x)$ for all $x \in \mathbb{R}$. $\gamma(t)$ is a moving boundary which keeps the mass of $f(t, \cdot)$ at 1, but the speed at which it moves is not known in advance and depends in turn on f .

It is not a priori obvious that (FB) has a solution, let alone that such a solution is unique. We will prove the existence and uniqueness using arguments closely related to the ones we will use to prove the existence of the limiting density.

We will denote by $\mathcal{P}$ the space of probability measures on $\mathbb{R}$, which we endow with the topology of weak convergence, and by $D([0, T], \mathcal{P})$ the space of càdlàg functions from $[0, T]$ to $\mathcal{P}$ endowed with the Skorohod topology.

Theorem 1. *For any fixed $T > 0$ the sequence of $\mathcal{P}$ -valued processes ν_t^N on $[0, T]$ converges in distribution in $D([0, T], \mathcal{P})$ to a deterministic $\nu_t \in D([0, T], \mathcal{P})$. ν_t is absolutely continuous with respect to the Lebesgue measure for every $t \in [0, T]$ and the corresponding densities $f(t, \cdot)$ are characterized by the following: there exists a continuous non-decreasing function $\gamma: [0, \infty) \rightarrow \mathbb{R}$ with $\gamma(0) = 0$ such that $(f(t, x), \gamma(t))$ is the unique solution of the free boundary problem (FB). In particular for $x > \gamma(t)$, $f(t, x)$ is strictly positive, jointly continuous in t and x and differentiable in t .*

Let us remark that there are at least two other examples in the literature of particle systems converging to the solution of a free boundary equation, but in both cases the limiting equation is of a different type. Chayes and Swindle (1996) study a particle system corresponding to a liquid-solid system with an interface subject to melting and freezing, while Gravner and Quastel (2000) study an internal diffusion limited aggregation model. In both cases an hydrodynamic limit is proved with the limiting equation corresponding to the famous Stefan problem, which involves free boundary problems for the heat equation where the moving boundary separates the solid and liquid phases (see Meirmanov (1992) and references therein for more on this problem).

1.2 Behavior of the finite system

To study the finite system it will be useful to introduce the *shifted* process Δ_t^N , which we define as follows:

$$\Delta_t^N = (\Delta_t^N(1), \dots, \Delta_t^N(N)) \quad \text{with} \quad \Delta_t^N(j) = \eta_t^N(j) - \eta_t^N(N).$$

Observe that $\Delta_t^N(N)$ is always 0. It is clear that Δ_t^N is also a Markov process, and its transitions are the same as those of η_t^N except that after erasing the leftmost particle

the N remaining particles are shifted to the left so that the new leftmost one lies at the origin.

We will denote by $\min \eta_t^N = \eta_t^N(N)$ and $\max \eta_t^N = \eta_t^N(1)$ the locations of the leftmost and rightmost particles in η_t^N .

Theorem 2. *For every fixed $N > 0$ the following hold:*

(a) *There is an $a_N > 0$ such that*

$$\lim_{t \rightarrow \infty} \frac{\min \eta_t^N}{t} = \lim_{t \rightarrow \infty} \frac{\max \eta_t^N}{t} = a_N$$

almost surely and in L^1 . Moreover, the sequence $(a_N)_{N>0}$ is non-decreasing.

(b) *The process Δ_t^N has a unique stationary distribution μ_N , which is absolutely continuous.*

(c) *For any (random or deterministic) initial condition ν_0 we have*

$$\|\mathbb{P}^{\nu_0}(\Delta_t^N \in \cdot) - \mu_N(\cdot)\|_{TV} \xrightarrow{t \rightarrow \infty} 0.$$

From this point on we will assume that the displacement distribution ρ has exponential decay. To be precise, we assume that there is an $\alpha > 0$ such that

$$(1.2) \quad \rho(x) \leq C e^{-\alpha|x|}.$$

for some $C > 0$. We will write

$$\Theta = \sup \left\{ \alpha > 0 : \sup_{x \in \mathbb{R}} [e^{\alpha|x|} \rho(x)] < \infty \right\}.$$

That is, $\Theta \in (0, \infty]$ is the maximal exponential rate of decay of ρ in the sense that $\rho(x) \leq C e^{-\alpha x}$ for some $C > 0$ when $\alpha < \Theta$ but not when $\alpha > \Theta$. Θ may be ∞ (as in the cases where ρ has compact support or ρ corresponds to a normal distribution), while $\Theta > 0$ is ensured by (1.2).

Now let $\phi(\theta) = \int_{-\infty}^{\infty} e^{\theta x} \rho(x) dx$ be the moment generating function of the displacement distribution ρ . (1.2) and our definition of Θ imply that $\phi(\theta) < \infty$ for $\theta \in (-\Theta, \Theta)$. To avoid unnecessary technical complications we will make the following extra assumption, which in particular implies that $\phi(\theta) = \infty$ for $|\theta| > \Theta$:

$$(1.3) \quad \frac{\phi(\theta)}{\theta} \xrightarrow{\theta \rightarrow \Theta^-} \infty.$$

This assumption always holds when $\Theta = \infty$: choosing $0 < l_1 < l_2$ so that $\rho(x) \geq M$ for some $M > 0$ and all $x \in [l_1, l_2]$ we get

$$\frac{1}{\theta} \int_{-\infty}^{\infty} e^{\theta x} \rho(x) dx \geq \frac{M(l_2 - l_1)}{\theta} e^{\theta l_1} \xrightarrow{\theta \rightarrow \infty} \infty.$$

Our next result will relate the asymptotic propagation speed a_N of our process η_t^N with the asymptotic speed of the rightmost particle in the branching random walk ξ_t^N .

Theorem 3.

$$\lim_{N \rightarrow \infty} a_N = a,$$

where a is the asymptotic speed of the rightmost particle in ξ_t^N , i.e., in a branching random walk where particles branch at rate 1 and their offspring are displaced by an amount chosen according to ρ .

Remark. If ρ is uniform on $[-1, 1]$ this follows from (8) in Durrett and Mayberry (2008). However, the couplings on which the proof are based extend easily to our more general setting, so we do not give the details of the proof.

The speed a has an explicit expression (see Biggins (1977)): by standard results of the theory of large deviations, if S_t is a continuous time random walk jumping at rate 1 and with jump distribution ρ , then the limit

$$(1.4) \quad \Lambda(x) = \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}(S_t > xt)$$

exists and equals $-(\sup_{\theta > 0} \{x\theta - \phi(\theta)\} + 1)$; a is given then by the formula $\Lambda(a) = -1$ (see Durrett and Mayberry (2008) for more on this).

1.3 Traveling wave solutions

A *traveling wave solution* of (FB) is a solution of the form $f(t, x) = w(x - ct)$ and $\gamma(t) = ct$ for some $c > 0$ (with initial condition $f_0(x) = w(x)$).

If w is a traveling wave solution, then from (FB1) we get

$$cw'(z) = - \int_0^\infty w(y)\rho(z - y) dy \quad \forall z > 0,$$

so integrating from $z = x$ to ∞ we deduce that w must solve

$$(1.5a) \quad w(x) = \frac{1}{c} \int_0^\infty w(y)R(x - y) dy \quad \forall x > 0,$$

$$(1.5b) \quad \int_0^\infty w(x) dx = 1,$$

$$(1.5c) \quad w(x) = 0 \quad \forall x \leq 0.$$

$$(1.5d) \quad w(x) \geq 0 \quad \forall x > 0,$$

where

$$R(x) = \int_x^\infty \rho(y) dy$$

is the tail distribution of ρ . On the other hand, it is easy to check that if w satisfies (1.5) and $f_0(x) = w(x)$, then $(w(x - ct), ct)$ is the solution of (FB).

(1.5a) is known as a Wiener-Hopf equation. Equations of this type have been studied since at least the 1920's (at the time in relation with the theory of radiative energy transfer), and have since been extensively studied and found relevance in diverse problems in mathematical physics and probability. In general, these equations can be solved using the Wiener-Hopf method, which was introduced in Wiener and Hopf (1931) (see

Chapter 4 of Paley and Wiener (1987) and also Kreĭn (1962)). But the solutions provided by this method are not necessarily positive, so they are not useful in our setting. Instead, we will rely on the results of Spitzer (1957), who studied these equations via probabilistic methods in the case where $R(x)/c$ is a probability kernel.

To do that we need to convert our equation to one where the kernel with respect to which we integrate is a probability kernel. To that end, we need to make the following observation. In Lemma 4.1 we will show that there is a $\lambda^* \in (0, \Theta)$ such that

$$(1.6) \quad \frac{\phi(\lambda^*)}{\lambda^*} = \min_{\lambda \in (0, \Theta)} \frac{\phi(\lambda)}{\lambda} = a,$$

where a is the asymptotic speed introduced in Theorem 3. On the other hand, the function $\lambda \mapsto \phi(\lambda)/\lambda$ is continuous and goes to ∞ as $\lambda \rightarrow 0$ (and moreover, as we will show in Lemma 4.1, it is decreasing on $(0, \lambda^*)$). Thus for every $c \geq a$ there is a $\lambda \in (0, \lambda^*]$ such that $\phi(\lambda)/\lambda = c$.

Observe that the tail distribution R of ρ has the same decay as ρ (see (1.2)). That is, for every $0 < \alpha < \Theta$ there is a $C > 0$ so that

$$(1.7) \quad R(x) \leq C e^{-\alpha x} \quad \forall x \geq 0.$$

Fix $c \geq a$ and use the above observation to pick a $\lambda \in (0, \lambda^*]$ such that $\phi(\lambda)/\lambda = c$. (1.7) implies that the function $x \mapsto e^{\lambda x} R(x)$ is in $L^1(\mathbb{R})$. Moreover, integration by parts yields

$$\int_{-\infty}^{\infty} e^{\lambda x} R(x) dx = \frac{\phi(\lambda)}{\lambda}.$$

Therefore

$$k(x) = \frac{\lambda}{\phi(\lambda)} e^{\lambda x} R(x)$$

is a probability kernel. On the other hand, if w is a solution of (1.5) with $c = \phi(\lambda)/\lambda$ then it is easy to check that $u(x) = e^{\lambda x} w(x)$ satisfies

$$(1.8) \quad u(x) = \int_0^{\infty} u(y) k(x-y) dy \quad \forall x \geq 0.$$

Thus the idea will be to recover solutions of (1.5) from positive solutions of (1.8).

Positive solutions of (1.8) can be regarded as densities of stationary measures for the following Markov chain. Let ξ_n be a sequence of i.i.d. random variables with distribution given by k , let $X_0 = 0$ and define

$$X_{n+1} = (X_n + \xi_n)^+,$$

where $x^+ = \max\{0, x\}$. This chain appears, for example, in the study of ladder variables for a random walk (see Chapter XII of Feller (1971)) and in the study of the GI/G/1 queue (see Chapter 5 of Durrett (2004)). If u satisfies (1.8), it is 0 on the negative half-line and it is non-negative on the positive half-line, then the measure (supported on $[0, \infty)$) having u as its density is invariant for X_n . Assuming that $\mathbb{E}(|\xi_1|) < \infty$, X_n is recurrent, null-recurrent or transient according to whether $\mathbb{E}(\xi_1)$ is negative, zero or positive. As we will see in Section 4, this expectation is negative in our case if and only

if $\lambda < \lambda^*$ and it is zero for $\lambda = \lambda^*$. In both cases the theory of recurrent Harris chains suggests (and Theorem 4.2 will prove) that there exists a unique (up to multiplicative constant) invariant measure for X_n , although this measure may not be finite in the null-recurrent case. The difference between the recurrent and null-recurrent cases explains the difference between the cases $c > a$ and $c = a$ in Theorem 4 below. The fact that the chain is transient when $\lambda > \lambda^*$ suggests that there are no positive solutions of (1.8) for these values of λ (again see Theorem 4.2 for a proof). This in turn hints at the possibility that there are no solutions of (1.5) for $c < a$. Our proof of this fact will not rely in seeing (1.5) as a Wiener-Hopf equation, but instead will use explicitly the fact that its solutions are traveling wave solutions of (FB).

Theorem 4. *Assume that (1.2) and (1.3) hold.*

- (a) *If $c \geq a$ the equation (1.5) has a unique solution w . This solution is differentiable except at the origin.*
- (b) *When $c > a$, and letting $\lambda \in (0, \lambda^*)$ be such that $\phi(\lambda)/\lambda = c$, the solution satisfies $\int_0^\infty e^{\lambda x} w(x) dx < \infty$ (which, in particular, implies that $\int_x^\infty w(y) dy = O(e^{-\lambda x})$). Moreover, if $\tilde{\lambda} > \lambda$ then $\sup_x e^{\tilde{\lambda} x} \int_x^\infty w(y) dy = \infty$.*
- (c) *When $c = a$ the solution satisfies $\int_0^x e^{\lambda^* y} w(y) dy = O(x)$.*
- (d) *If $c < a$ the equation (1.5) has no solution.*

We remark that, when $c > a$, the solution w given by the theorem can be obtained by the following limiting procedure. Take $0 < \lambda < \lambda^*$ as in the above statement and let u_0 be the density of any non-negative random variable whose distribution is absolutely continuous. Now let $w_0(x) = e^{-\lambda x} u_0(x)$ and then for $n \geq 1$ let

$$w_{n+1}(x) = \frac{1}{c} \int_0^\infty w_n(y) R(x-y) dy \quad \text{for } x \geq 0.$$

Then the limit $w_\infty(x) = \lim_{n \rightarrow \infty} w_n(x)$ exists and defines an integrable continuous function. The solution w is then given by $w(x) = K w_\infty(x)$ with $K > 0$ chosen so that w integrates to 1. The fact that w has this representation follows from the results of Spitzer (see Theorem 4.2).

The rest of the paper is devoted to proofs, with one section devoted to each one of the proofs of Theorems 1, 2 and 4.

2 Proof of Theorem 1

Most of the work in the proof of Theorem 1 will correspond to showing that for each fixed $t \geq 0$ the tail distribution of our process at that time, $F^N(t, x) = \nu_t^N([x, \infty))$, converges (almost surely and in L^1) to a deterministic limit $F(t, x)$ corresponding to the tail distribution of a random variable and that, moreover, the limit $F(t, x)$ has a density $f(t, x)$ (that is $F(t, x) = \int_x^\infty f(t, y) dy$) which solves (FB).

To get started we need to introduce some auxiliary processes. Without loss of generality we may work during all this section with η_t^N defined on the time interval $[0, T]$ for some given T , and for simplicity we will actually assume that $T = 1$.

First, recall that ξ_t^N is the same process as η_t^N but with no killing and let $\hat{\nu}_t^N$ be the associated empirical measure. Let $\mathcal{M}$ be the space of finite measures on $\mathbb{R}$, endowed

with the topology of weak convergence, and let $C([0, 1], \mathcal{M})$ and $D([0, 1], \mathcal{M})$ be the spaces of continuous and càdlàg functions from $[0, 1]$ to $\mathcal{M}$ endowed respectively with the uniform and Skorohod topologies.

Proposition 2.1. *Fix $T > 0$. Then the process $\widehat{\nu}_t^N$ converges in distribution in $D([0, 1], \mathcal{P})$ to a deterministic $\widehat{\nu}_t$ in $C([0, 1], \mathcal{P})$ which for each $t \in [0, 1]$ is absolutely continuous with respect to the Lebesgue measure. If we denote the density of $\widehat{\nu}_t$ by $\widehat{f}(t, x)$ then $\widehat{f}(t, x)$ is the unique solution to the integro-differential equation*

$$(2.1) \quad \frac{\partial \widehat{f}}{\partial t}(t, x) = \int_{-\infty}^{\infty} \widehat{f}(t, y) \rho(x - y) dy$$

on $[0, 1]$ with initial condition $\widehat{f}(0, x) = f_0(x)$.

Proof. By Theorem 5.3 of Fournier and Méléard (2004) we have that $\widehat{\nu}_t^N$ converges in distribution to a deterministic $\widehat{\nu}_t$ in $C([0, 1], \mathcal{P})$ which is the unique solution of the following system: for all bounded and measurable φ ,

$$(2.2) \quad \int_{-\infty}^{\infty} \varphi(x) \widehat{\nu}_t(dx) = \int_{-\infty}^{\infty} \varphi(x) \widehat{\nu}_0(dx) + \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \varphi(y) \rho(x - y) dy \widehat{\nu}_s(dx) ds.$$

Moreover, by Proposition 5.4 of the same paper, $\widehat{\nu}_t$ is absolutely continuous for all $t \in [0, 1]$, and hence its density $\widehat{f}(t, x)$ must satisfy $\widehat{f}(0, x) = f_0(x)$ and

$$\frac{d}{dt} \int_{-\infty}^{\infty} \varphi(x) \widehat{f}(t, x) dx = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \varphi(y) \rho(x - y) \widehat{f}(t, x) dy dx.$$

Taking $\varphi = \mathbf{1}_{[z, z+h]}$, dividing by h , taking $h \rightarrow 0$ and using the symmetry of ρ we deduce that $\widehat{f}$ satisfies (FB1) at z and the result follows. $\square$

Recalling that $f_0(x) > 0$ for $x \geq 0$ it is clear that if $\widehat{f}$ solves (2.1) then $\widehat{f}(t, x) > 0$ for all $t \in [0, 1]$ and $x \in \mathbb{R}$.

For each $k \geq 0$ we define the modified process $\eta_t^{N,k}$ inductively as follows. For each $m = 0, \dots, 2^k - 1$, run the process with no killing on the interval $[\frac{m}{2^k}, \frac{m+1}{2^k})$ and then at time $\frac{m+1}{2^k}$ repeatedly delete the leftmost particle until there are only N left. Let $\nu_t^{N,k}$ be the empirical measure associated with this process (we consider here the càdlàg version of this process, and we do the same for other processes and functions defined below). To make clear the distinction between the particles we erase in η_t^N and those we erase in this modified process at dyadic times, we will refer to this last procedure as *shaving off the extra mass* in $\eta_t^{N,k}$. Our goal now is to study the convergence of the tail distribution of $\nu_t^{N,k}$.

Let us first describe what will turn out to be the limit. It is defined inductively on each of the dyadic subintervals of $[0, 1]$. We let $f^k(0, x) = f_0(x)$. If we have constructed f^k up to time $\frac{m}{2^k}$ for some $m \in \{0, \dots, 2^k - 1\}$ then for $t \in (\frac{m}{2^k}, \frac{m+1}{2^k})$ we let $f^k(t, x)$ be the solution of

$$(2.3) \quad \frac{\partial f^k}{\partial t}(t, x) = \int_{-\infty}^{\infty} f^k(t, y) \rho(x - y) dy.$$

Then at time $\frac{m+1}{2^k}$ we let X_{m+1}^k be such that

$$\int_{X_{m+1}^k}^{\infty} f^k\left(\left(\frac{m+1}{2^k}\right)-, y\right) dy = 1$$

and define

$$f^k\left(\frac{m+1}{2^k}, x\right) = f^k\left(\left(\frac{m+1}{2^k}\right)-, x\right) \mathbf{1}_{x > X_{m+1}^k}.$$

We will denote by $F^{N,k}$ the tail distribution of $\nu_t^{N,k}$, that is,

$$F^{N,k}(t, x) = \nu_t^{N,k}([x, \infty)),$$

and by F^k be the tail distribution of ν_t^k .

Remark. To avoid confusion (and notational complications) we will use the following convention: upper-case superscripts, such as in $F^N(t, x)$, refer to quantities associated with our stochastic process η_t^N , while lower-case superscripts, such as in $F^k(t, x)$ refer to the deterministic quantities we just introduced.

Proposition 2.2. *For every fixed $k \geq 1$ and every $t \in [0, 1]$ and $x \in \mathbb{R}$, $F^{N,k}(t, x)$ converges in probability to $F^k(t, x)$.*

Proof. Proposition 2.1 implies that $\nu_t^{N,k}([x, \infty))$ converges in probability to $\nu_t^k([x, \infty))$ for all $t \in [0, \frac{1}{2^k})$ and all $x \in \mathbb{R}$. Using the partial order in $\mathcal{M}$ given by $\mu \preceq \nu$ if and only if $\mu([x, \infty)) \leq \nu([x, \infty))$ for all $x \in \mathbb{R}$, it is clear that the mappings $t \mapsto \nu_t^k$ and $t \mapsto \nu_t^{N,k}$ are increasing on $[0, \frac{1}{2^k})$, and thus the limits $\lim_{t \uparrow 1/2^k} \nu_t^k = \nu_-^k$ and $\lim_{t \uparrow 1/2^k} \nu_t^{N,k} = \nu_-^{N,k}$ exist (for $\nu_t^{N,k}$ these statements hold almost surely). Using the inequalities

$$\begin{aligned} (\nu_t^k - \nu_s^k)(\mathbb{R}) &\leq e^t - e^s \quad \text{and} \\ \mathbb{E}\left((\nu_t^{N,k} - \nu_s^{N,k})(\mathbb{R}) \middle| \nu_s^{N,k}\right) &\leq e^{t-s} \nu_s^{N,k}(\mathbb{R}) \end{aligned}$$

(which follow from the comment after (1.1)) and the fact that ν_-^k is absolutely continuous, it is easy to see that $\nu_-^{N,k}([x, \infty))$ converges in probability to $\nu_-^k([x, \infty))$ for all x . Since $x \mapsto \nu_-^k([x, \infty))$ is strictly decreasing at $x = X_1^k$, the location of the point at which the mass of $\nu_t^{N,k}$ is shaved off at time $\frac{1}{2^k}$ converges in probability to X_1^k . The result for $t = \frac{1}{2^k}$ follows from this, and induction gives the desired result. $\square$

Proposition 2.3. *For every given $t \in [0, 1]$ and $x \in \mathbb{R}$, $F^{N,k}(t, x)$ converges in probability to $F^N(t, x)$ as $k \rightarrow \infty$, uniformly in N .*

The proof depends on the following lemma:

Lemma 2.4. *We can couple $\eta_t^{N,k}$ and η_t^N (starting with the same initial configuration) in such a way that the following holds: for every $N \geq 1$, $k \geq 1$ and $m \in \{0, \dots, 2^k\}$,*

$$\mathbb{E}(|\eta_{m/2^k}^N \Delta \eta_{m/2^k}^{N,k}|) \leq 2N \frac{e-1}{e^{2^{-k}}-1} 2^{-2k},$$

where $A \Delta B = A \setminus B \cup B \setminus A$.

Proof. The coupling will be constructed inductively on each dyadic subinterval of $[0, 1]$. We start both processes with the same initial configuration. The idea will be to use the same branching times and displacements whenever possible. To do this we will decompose $\eta_t^{N,k}$ in the following way

$$(2.4) \quad \eta_t^{N,k} = G_t^{N,k} \cup D_t^{N,k} \cup B_t^{N,k},$$

where the unions are disjoint, $G_t^{N,k} \subseteq \eta_t^N$ will correspond to “good particles” (i.e. coupled particles), $B_t^{N,k}$ will correspond to “bad particles” (i.e. uncoupled particles) and $D_t^{N,k}$ will correspond to “dangerous particles” (i.e. particles which might become bad if not erased).

To get started we take $G_0^{N,k} = \eta_0^N$ and $D_0^{N,k} = B_0^{N,k} = \emptyset$. Let $0 \leq \tau_1 < \tau_2 < \dots \leq 1$ be the sequence of branching times for particles in η_t^N (there are almost surely finitely many such times) and let $I_m \in \mathbb{N}$ be such that τ_{I_m} is the last branching before time $\frac{m}{2^k}$. For each $i \leq I_1$ we check whether the branching at time $\tau_i \leq \frac{1}{2^k}$ occurs at a particle which is in $G_{\tau_i}^{N,k}$. If that is the case, we add the new particle to $G_{\tau_i}^{N,k}$ and then we look for the particle erased from $\eta_{\tau_i}^N$ after this branching and move it from $G_t^{N,k}$ to $D_t^{N,k}$. By our decomposition this results in $\eta_t^{N,k}$ having a branching at each time τ_i and using the same displacements as in η_t^N to choose the location of the newborn particles. The difference is that in η_t^N the leftmost particle is erased, while in $\eta_t^{N,k}$ the leftmost particle is moved to the set of dangerous particles. At the same time, we let each particle in $D_t^{N,k} \cup B_t^{N,k}$ branch at rate 1 and use ρ to choose the displacements for their offspring. Whenever there is such a branching we put the newborn particle in $B_t^{N,k}$. In this manner, every particle that is in $\eta_t^{N,k} \cap \eta_t^N$ undergoes branchings at the same times and uses the same displacements for the newborn particles, while the particles in $\eta_t^{N,k} \setminus \eta_t^N$ branch independently and all their offspring is labelled as bad. At time $\frac{1}{2^k}$ we need to shave off the extra mass in $\eta_t^{N,k}$. Observe that, by our construction, all dangerous particles are erased (though we will, in general, also erase good and bad particles).

We continue the coupling inductively following the same procedure. For $1 \leq m \leq 2^k - 1$ we will have a decomposition $\eta_{m/2^k}^{N,k} = G_{m/2^k}^{N,k} \cup D_{m/2^k}^{N,k} \cup B_{m/2^k}^{N,k}$ (in fact, $D_{m/2^k}^{N,k}$ will be empty, as we already mentioned for the case $m = 1$, and by the discussion below for $m > 1$). We consider the branching times τ_i for $I_m < i \leq I_{m+1}$ and use these for the shared branchings (i.e., for those particles in $G_t^{N,k}$) on the time interval $(\frac{m}{2^k}, \frac{m+1}{2^k})$. Whenever the erased particle in η_t^N is also in $G_t^{N,k}$, we move it from that set to $D_t^{N,k}$. We keep branching the remaining particles (those in $\eta_t^N \Delta \eta_t^{N,k}$) independently as described above. At time $\frac{m+1}{2^k}$ we need to shave off the extra mass in $\eta_t^{N,k}$. As before, we may erase particles of each of the three types, but it is no longer necessarily the case that all dangerous particles will be erased. In any case, the total number of dangerous and bad particles remaining must be at most the number of bad particles before shaving. To see this, observe that each dangerous particle corresponds to a branching which increased by one the number of particles in $\eta_t^{N,k}$, so to each dangerous particle there corresponds some particle which will be erased when shaving; the corresponding particle to be erased is possibly the dangerous particle itself, and otherwise it has to be a bad particle because all good particles are to the right of any dangerous particle. After shaving off the mass we relabel all the remaining dangerous particles as bad.

An important consequence of the preceding is that the sequence $(|B_{m/2^k}^{N,k}|)_{m=0,\dots,2^k}$ is non-decreasing (because the total number of particles at these times is always N and the number of dangerous particles at these times is 0, while the number of good particles cannot grow).

On each dyadic subinterval the pair $(|D_t^{N,k}|, |B_t^{N,k}|)$ is dominated by a process (d_t^k, b_t^k) which evolves as follows:

$$\begin{array}{lll} d_t^k & \longrightarrow & d_t^k + 1 & \text{at rate } N \\ b_t^k & \longrightarrow & b_t^k + 1 & \text{at rate } d_t^k + b_t^k \end{array}$$

(the second rate is the correct one, but the first one is not necessarily so, because the number of good particles is in general less than or equal to N). Using the fact that d_t^k is always 0 at the beginning of each dyadic subinterval, this implies that $\mathbb{E}(b_t^k)$ must solve $d\mathbb{E}(b_t^k)/dt = Nt + \mathbb{E}(b_t^k)$ on these intervals. The solution of the ODE $y'(t) = Nt + y(t)$ is $y(t) = (y(0) + N)e^t - N(1 + t)$, and thus for $\frac{m}{2^k} \leq t < \frac{m+1}{2^k}$ we obtain

$$(2.5) \quad \mathbb{E}(b_t^k) = (\mathbb{E}(b_{m/2^k}^k) + N)e^{t - \frac{m}{2^k}} - N(t - \frac{m}{2^k} + 1).$$

Since $b_0^k = 0$ we deduce that $\mathbb{E}(b_{(1/2^k)_-}^k) = N(e^{2^{-k}} - 2^{-k} - 1) \leq N2^{-2k}$. At time $\frac{1}{2^k}$ we need to shave off the extra mass in $\eta_{(1/2^k)_-}^{N,k}$ and this leaves us with $d_{1/2^k}^k = 0$ and $b_{1/2^k}^k \leq b_{(1/2^k)_-}^k$. Repeating this argument we get

$$\mathbb{E}(b_{2/2^k}^k) \leq \mathbb{E}(b_{(2/2^k)_-}^k) = N[(1 + 2^{-2k})e^{2^{-k}} - (2^{-k} + 1)] \leq N(1 + e^{2^{-k}})2^{-2k}$$

and inductively we deduce that

$$(2.6) \quad \mathbb{E}(b_1^k) \leq N \left[\sum_{j=0}^{2^k-1} e^{j2^{-k}} \right] 2^{-2k} = N \frac{1 - e}{1 - e^{2^{-k}}} 2^{-2k}.$$

Since by our construction there are no dangerous particles at time 1 we have $|\eta_1^N \Delta \eta_1^{N,k}| = 2|B_1^{N,k}|$ and the result follows for $m = 2^k$. Since $(|B_{m/2^k}^{N,k}|)_{m=0,\dots,2^k}$ is non-decreasing, the same holds for $m < 2^k$. $\square$

Proof of Proposition 2.3. Fix $k > 0$ for a moment and assume that $t \in [\frac{m}{2^k}, \frac{m+1}{2^k})$. Using the coupling introduced in the proof Lemma 2.4 we have that $\mathbb{E}(|B_{m/2^k}^{N,k}|) < N2^{-2k}(e - 1)/(e^{2^{-k}} - 1)$ and thus (2.5) implies that

$$\begin{aligned} \mathbb{E}(|\eta_t^N \Delta \eta_t^{N,k}|) &= 2\mathbb{E}(|D_t^{N,k}|) + 2\mathbb{E}(|B_t^{N,k}|) \\ &\leq 2N(t - \frac{m}{2^k}) + 2N \left[\left(1 + \frac{e - 1}{e^{2^{-k}} - 1} 2^{-2k}\right) e^{t - m/2^k} - (t - \frac{m}{2^k}) - 1 \right] \\ &\leq N2^{-k+1} + N \left[\frac{e - 1}{e^{2^{-k}} - 1} e^{2^{-k}} + 1 \right] 2^{-2k+1}. \end{aligned}$$

Observing that

$$\left| F^{N,k}(t, x) - F^N(t, x) \right| = \frac{1}{N} \left| \sum_{i=1}^N \left(\mathbf{1}_{\eta_i^{N,k}(i) \geq x} - \mathbf{1}_{\eta_i^N(i) \geq x} \right) \right| \leq \frac{1}{N} \sum_{i=1}^N \mathbf{1}_{\eta_i^{N,k}(i) \neq \eta_i^N(i)}$$

we deduce by Markov's inequality and the above estimate that

$$\mathbb{P}\left(\left|F^{N,k}(t, x) - F^N(t, x)\right| > \varepsilon\right) \leq \frac{1}{\varepsilon} \left[2^{-k+1} + \left(\frac{e-1}{e^{2^{-k}}-1} e^{2^{-k}} + 1\right) 2^{-2k+1}\right] \xrightarrow[k \rightarrow \infty]{} 0$$

uniformly in N . $\square$

Lemma 2.5.

$$F^k(t, x) \geq F^{k+1}(t, x) \quad \text{for all } t \in [0, 1], x \in \mathbb{R}, k \geq 1.$$

Proof. Fix $k \geq 1$ and $x \in \mathbb{R}$. The result is trivial at $t = 0$. We will work inductively on the intervals $(\frac{m}{2^k}, \frac{m+1}{2^k}]$. Take $0 \leq m \leq 2^k - 1$ and assume that

$$F^k(\frac{m}{2^k}, x) \geq F^{k+1}(\frac{m}{2^k}, x).$$

Then writing $H = F^k - F^{k+1}$ we have for $t \in (\frac{m}{2^k}, \frac{m+1}{2^k})$ that

$$\begin{aligned} \frac{\partial H}{\partial t}(t, x) &= \int_x^\infty \int_{-\infty}^\infty [f^k(t, u) - f^{k+1}(t, u)] \rho(y - u) du dy \\ (2.7) \quad &= \int_{-\infty}^\infty \int_x^\infty [f^k(t, y - v) - f^{k+1}(t, y - v)] \rho(v) dy dv \\ &= \int_{-\infty}^\infty H(t, x - v) \rho(v) dv = \int_{-\infty}^\infty H(t, z) \rho(x - z) dz, \end{aligned}$$

so H satisfies (FB1) on this interval and thus, since $H(\frac{m}{2^k}, \cdot) \geq 0$, we get $H(t, \cdot) \geq 0$ for $t \in (\frac{m}{2^k}, \frac{m+1}{2^k})$. At time $\frac{2m+1}{2^{k+1}} = \frac{m}{2^k} + \frac{1}{2^{k+1}}$ the density f^{k+1} is shaved off, leaving $F^k(\frac{2m+1}{2^{k+1}}, x) \geq F^{k+1}(\frac{2m+1}{2^{k+1}}, x)$. Repeating the above argument we get

$$(2.8) \quad F^k(t, x) \geq F^{k+1}(t, x) \quad \text{for all } t \in (\frac{m}{2^{k+1}}, \frac{m+1}{2^{k+1}}).$$

Now at time $\frac{m+1}{2^{k+1}}$ both densities f^k and f^{k+1} are shaved off, say at points x_k and x_{k+1} , respectively. Then by (2.8), $x_k \geq x_{k+1}$, and thus (2.8) holds at $t = \frac{m+1}{2^{k+1}}$ as well. $\square$

Since $F^k(t, x)$ is decreasing and positive, we can define

$$F(t, x) = \lim_{k \rightarrow \infty} F^k(t, x).$$

It is obvious that for each given t , $F(t, \cdot)$ is non-increasing and its range is $[0, 1]$.

Proposition 2.6. *For every $t \in [0, 1]$ and $x \in \mathbb{R}$, $F^N(t, x)$ converges almost surely and in L^1 as $N \rightarrow \infty$ to $F(t, x)$.*

Proof. First observe that, for fixed $t \in [0, 1]$ and $x \in \mathbb{R}$ and since $F^N(t, x) \leq 1$, the sequence of random variables $(F^N(t, x))_{N \geq 0}$ is uniformly integrable, so it is enough to show that $F^N(t, x) \rightarrow F(t, x)$ in probability.

Fix $\varepsilon > 0$. Use Proposition 2.2 to choose, for each $k > 0$, an $N_k > 0$ so that

$$\mathbb{P}\left(\left|F^{N,k}(t, x) - F^k(t, x)\right| > \frac{\varepsilon}{2}\right) < \frac{1}{k}$$

for every $N \geq N_k$ and $N_k \uparrow \infty$. Define k_N as follows: $k_N = 1$ for $N < N_1$ and $k_N = k$ for $N_k \leq N \leq N_{k+1}$. We have

$$\begin{aligned} \mathbb{P}\left(\left|F^{N,k_N}(t,x) - F(t,x)\right| > \varepsilon\right) &\leq \mathbb{P}\left(\left|F^{N,k_N}(t,x) - F^{k_N}(t,x)\right| > \frac{\varepsilon}{2}\right) \\ &\quad + \mathbb{P}\left(\left|F^{k_N}(t,x) - F(t,x)\right| > \frac{\varepsilon}{2}\right). \end{aligned}$$

By the definition of k_N , the first term on the right hand side is less than $1/k_N$, while the second one is 0 when k_N is large enough. We deduce that $F^{N,k_N}(t,x)$ converges in probability as $N \rightarrow \infty$ to $F(t,x)$.

To finish the proof write

$$\begin{aligned} \mathbb{P}\left(\left|F^N(t,x) - F(t,x)\right| > \varepsilon\right) &\leq \mathbb{P}\left(\left|F^N(t,x) - F^{N,k_N}(t,x)\right| > \frac{\varepsilon}{2}\right) \\ &\quad + \mathbb{P}\left(\left|F^{N,k_N}(t,x) - F(t,x)\right| > \frac{\varepsilon}{2}\right). \end{aligned}$$

We already know that the second term on the right hand side goes to 0, while the first one goes to 0 thanks to Proposition 2.3 (here we use the fact that the convergence is uniform in N). $\square$

Recall the definition of the shaving points X_m^k and let $X^k : [0,1] \rightarrow \mathbb{R}$ be the corresponding linear interpolation, that is,

$$X^k(t) = X_m^k + \frac{X_{m+1}^k - X_m^k}{2^k} \left(t - \frac{m}{2^k}\right) \quad \text{for } \frac{m}{2^k} < t \leq \frac{m+1}{2^k}.$$

Lemma 2.7. $X^k(t)$ converges uniformly in $[0,1]$ to a continuous function $\gamma(t)$.

Proof. We will start by showing that the sequence of functions $(X^k)_{k>0}$ is relatively compact. By the Arzelà-Ascoli Theorem, we only need to show that our sequence is uniformly bounded and equicontinuous.

Observe that, for each given k , $X^k(t)$ is increasing. Indeed, it is enough to show that $X_m^k \leq X_{m+1}^k$ for $0 \leq m < 2^k$, and this follows from the fact that, if $f^k(\frac{m}{2^k}, \cdot) \geq 0$, then (FB1) implies that $f^k(t, \cdot) \geq f^k(\frac{m}{2^k}, \cdot)$ for $\frac{m}{2^k} < t < \frac{m+1}{2^k}$. Therefore

$$\sup_{k>0} \sup_{t \in [0,1]} X^k(t) = \sup_{k>0} X^k(1).$$

To show that this last supremum is finite, observe that $\widehat{f}(t,x)$ (which was defined in Proposition 2.1) satisfies $\widehat{f}(t,x) \geq f^k(t,x)$ for all k . On the other hand,

$$\int_{-\infty}^{\infty} \widehat{f}(t,x) dx = 1 + \int_0^t \int_{-\infty}^{\infty} \widehat{f}(s,x) dx ds$$

whence it is easy to see that

$$(2.9) \quad \int_{-\infty}^{\infty} \widehat{f}(t,x) dx = e^t.$$

Therefore, if $M > 0$ is such that $\int_M^{\infty} \widehat{f}(1,x) dx < 1$ we deduce that $X^k(1) \leq M$ for all k and the uniform boundedness follows.

For the equicontinuity we need to show that given any $\varepsilon > 0$ there is a $\delta > 0$ such that

$$\sup_{k>0} \left| X^k(t) - X^k(s) \right| < \varepsilon$$

whenever $|t - s| < \delta$. Assume that $s < t$, fix k for a moment and let $\frac{l}{2^k}$ and $\frac{m}{2^k}$ be the dyadic numbers immediately to the right of s and t , respectively (here we assume k is large enough so that $m \vee l < 2^k$). Then

$$(2.10) \quad \begin{aligned} \left| X^k(t) - X^k(s) \right| &\leq \left| X^k(t) - X^k\left(\frac{m}{2^k}\right) \right| + \left| X^k\left(\frac{m}{2^k}\right) - X^k\left(\frac{l}{2^k}\right) \right| + \left| X^k\left(\frac{l}{2^k}\right) - X^k(s) \right| \\ &\leq \left| X^k\left(\frac{m+1}{2^k}\right) - X^k\left(\frac{m}{2^k}\right) \right| + \left| X^k\left(\frac{m}{2^k}\right) - X^k\left(\frac{l}{2^k}\right) \right| + \left| X^k\left(\frac{l+1}{2^k}\right) - X^k\left(\frac{l}{2^k}\right) \right|. \end{aligned}$$

Now for any $p, q \in \{0, \dots, 2^k\}$ with $q \geq p$ we have

$$(2.11) \quad \begin{aligned} \int_{X^k(\frac{p}{2^k})+\varepsilon}^{\infty} f^k\left(\frac{q}{2^k}, y\right) dy &= \int_{X^k(\frac{p}{2^k})}^{\infty} f^k\left(\frac{p}{2^k}, y\right) dy + \int_{X^k(\frac{p}{2^k})}^{\infty} \left[f^k\left(\frac{q}{2^k}, y\right) - f^k\left(\frac{p}{2^k}, y\right) \right] dy \\ &\quad - \int_{X^k(\frac{p}{2^k})}^{X^k(\frac{p}{2^k})+\varepsilon} f^k\left(\frac{q}{2^k}, y\right) dy. \end{aligned}$$

The first term on the right hand side equals 1. The second term corresponds to the amount of mass accumulated by f^k to the right of $X^k(\frac{p}{2^k})$ on the time interval $(\frac{p}{2^k}, \frac{q}{2^k}]$. Using (FB1) it is not hard to see that this is bounded by the same quantity with f^k replaced by $\hat{f}$, which gives the bound

$$\int_{X^k(\frac{p}{2^k})}^{\infty} \left[\hat{f}\left(\frac{q}{2^k}, y\right) - \hat{f}\left(\frac{p}{2^k}, y\right) \right] dy \leq e^{\frac{q}{2^k}} - e^{\frac{p}{2^k}} \leq e^{\frac{q-p}{2^k}}$$

thanks to (2.9). On the other hand, using the fact that $X^k(t)$ is increasing, it is clear that $f^k(\frac{q}{2^k}, x) \geq f^k(0, x) = f_0(x)$ for $x \geq X^k(\frac{q}{2^k})$. Therefore the last term on (2.11) is greater than or equal to

$$\int_{X^k(\frac{p}{2^k})}^{X^k(\frac{p}{2^k})+\varepsilon} f_0(y) dy.$$

Now $X^k(\frac{p}{2^k})$ is non-negative and bounded by M by the preceding arguments, so the last integral is at least

$$L = \inf_{x \in [0, M]} \int_x^{x+\varepsilon} f_0(y) dy > 0$$

where we used the fact that f_0 is strictly positive on the positive half-line. Putting the last two bounds together with (2.11) we get

$$\int_{X^k(\frac{p}{2^k})+\varepsilon}^{\infty} f^k\left(\frac{q}{2^k}, y\right) dy \leq 1 + e^{\frac{q-p}{2^k}} - L.$$

Now $|t - s| < \delta$ implies that $\frac{m-l}{2^k} \leq \delta + \frac{1}{2^k}$, and thus we deduce that

$$\int_{X^k(\frac{p}{2^k})+\varepsilon}^{\infty} f^k\left(\frac{q}{2^k}, y\right) dy < 1$$

for small enough δ and large enough k and for $(p, q) \in \{(l, l+1), (m, m+1), (l, m)\}$. The preceding means that if δ is small enough and K is large enough then $|X^k(\frac{q}{2^k}) - X^k(\frac{p}{2^k})| < \varepsilon$ for $k \geq K$ and for these three pairs (p, q) . Using (2.10) we obtain

$$\sup_{k \geq K} |X^k(t) - X^k(s)| < \varepsilon$$

if $|t - s| < \delta$ and δ is small enough. Since the functions X^k are all uniformly continuous (on $[0, 1]$), it is clear that, by choosing δ even smaller if necessary, the same will hold also for $k = 1, \dots, K - 1$. This finishes the proof of the equicontinuity.

The last thing we need to show is that our sequence has a unique limit point. Consider two convergent subsequences $X^{n_k} \rightarrow \gamma_1$ and $X^{m_k} \rightarrow \gamma_2$. Let $t = \frac{i}{2^l}$ be any dyadic number in $[0, 1]$ and assume that k is large enough so that $n_k \wedge m_k \geq l$. Recall from the proof of Lemma 2.5 that $X^k(t)$ is non-increasing in k for each fixed $t \in [0, 1]$. Since $F^{n_k}(t, x) = 1$ for all $x \leq X^{n_k}(t)$ we deduce that

$$F^{n_k}(t, x) = 1 \quad \text{for all } x \leq \gamma_1(t).$$

Now given any k there is a k' such that $n_{k'} \geq m_k$, so by Lemma 2.5 we get

$$1 = F^{n_{k'}}(t, x) \leq F^{m_k}(t, x) \leq 1 \quad \text{for all } x \leq \gamma_1(t).$$

This means that $X^{m_k}(t) \geq \gamma_1(t)$ for all large enough k , and taking $k \rightarrow \infty$ we deduce that $\gamma_2(t) \geq \gamma_1(t)$. By symmetry we get $\gamma_1(t) \geq \gamma_2(t)$. This gives $\gamma_1(t) = \gamma_2(t)$ for all dyadic $t \in [0, 1]$, and now the uniqueness follows from the continuity of γ_1 and γ_2 . $\square$

Now we make the following observation. Suppose that $(g(t, x), \gamma(t))$ solves (FB) and let $G(t, x) = \int_x^\infty g(t, y) dy$. Then it is not difficult to check, repeating the arguments leading to (2.7), that $(G(t, x), \gamma(t))$ must solve the following free boundary problem (FB'):

$$(FB1) \quad \frac{\partial G}{\partial t}(t, x) = \int_{-\infty}^\infty G(t, y) \rho(x - y) dy \quad \forall x > \gamma(t)$$

$$(FB2') \quad G(t, x) = 1 \quad \forall x \leq \gamma(t)$$

with initial condition $G(0, x) = \int_x^\infty f_0(y) dy$. Moreover, if $(G(t, x), \gamma(t))$ solves (FB') and $G(t, \cdot)$ is absolutely continuous for all t , then $(g(t, x), \gamma(t))$, where $g(t, \cdot)$ is the density of $G(t, \cdot)$, must solve (FB).

Proposition 2.8. *$F(t, x)$ is differentiable in t for all $x > \gamma(t)$ and it satisfies (FB')*

Proof. We already proved that $(F(t, x), \gamma(t))$ satisfies (FB2'). For $x > \gamma(t)$ and by the definition of $F^k(t, x)$ (which implies that $F^k(t, x)$ is differentiable inside each dyadic

subinterval) we may write

$$\begin{aligned}
F^k(t, x) &= F^k(0, x) + \sum_{m=1}^{n_k(t)} \left[F^k\left(\frac{m}{2^k}, x\right) - F^k\left(\frac{m-1}{2^k}, x\right) \right] + \left[F^k(t, x) - F^k(n_k(t), x) \right], \\
&= F^k(0, x) + \sum_{m=1}^{n_k(t)} \left[\int_{(\frac{m-1}{2^k}, \frac{m}{2^k})} \frac{\partial F^k}{\partial s}(s, x) ds \right] + \int_{(\frac{n_k(t)}{2^k}, t)} \frac{\partial F^k}{\partial s}(s, x) ds \\
&\quad + \sum_{m=1}^{n_k(t)} \left[F^k\left(\frac{m}{2^k}, x\right) - F^k\left(\left(\frac{m}{2^k}\right)^-, x\right) \right] + \left[F^k(t, x) - F^k(t^-, x) \right],
\end{aligned}$$

where $n_k(t) = \lfloor \frac{2^k t}{1} \rfloor$. Recalling that $X^k(t) \downarrow \gamma(t)$, we can take k large enough so that $\gamma(t) \leq X^k(t) < x$. Since $X^k(s)$ is increasing in s we deduce that $X^k(\frac{m}{2^k}) < x$ for $m = 1, \dots, n_k(t)$, and therefore all the terms in the last line above are 0. On the other hand, observe that F^k must solve (FB1) on each dyadic subinterval, which can be checked repeating again the calculations in (2.7). Therefore,

$$\begin{aligned}
(2.12) \quad F^k(t, x) &= F^k(0, x) + \sum_{m=1}^{n_k(t)} \int_{(\frac{m-1}{2^k}, \frac{m}{2^k})} \int_{-\infty}^{\infty} F^k(s, y) \rho(x - y) dy ds \\
&\quad + \int_{(\frac{n_k(t)}{2^k}, t)} \int_{-\infty}^{\infty} F^k(s, y) \rho(x - y) dy ds \\
&= F^k(0, x) + \int_0^t \int_{-\infty}^{\infty} F(s, y) \rho(x - y) dy ds \\
&\quad + \sum_{m=1}^{n_k(t)} \int_{(\frac{m-1}{2^k}, \frac{m}{2^k})} \int_{-\infty}^{\infty} \left[F^k(s, y) - F(s, y) \right] \rho(x - y) dy ds \\
&\quad + \int_{(\frac{n_k(t)}{2^k}, t)} \int_{-\infty}^{\infty} \left[F^k(s, y) - F(s, y) \right] \rho(x - y) dy ds.
\end{aligned}$$

Now for fixed y , $F^k(\cdot, y)$ is a decreasing sequence converging to $F(\cdot, y)$, so Dini's Theorem implies that

$$\Delta_k(y) = \sup_{t \in [0, 1]} \left| F^k(t, y) - F(t, y) \right| \xrightarrow[k \rightarrow \infty]{} 0.$$

The sum of the terms on the last two lines of (2.12) is bounded by

$$\begin{aligned}
&\sum_{m=1}^{n_k(t)} \int_{(\frac{m-1}{2^k}, \frac{m}{2^k})} \int_{-\infty}^{\infty} \Delta_k(y) \rho(x - y) dy ds + \int_{(\frac{n_k(t)}{2^k}, t)} \int_{-\infty}^{\infty} \Delta_k(y) \rho(x - y) dy ds \\
&\leq t \int_{-\infty}^{\infty} \Delta_k(y) \rho(x - y) dy,
\end{aligned}$$

and this last integral goes to 0 as $k \rightarrow \infty$ by the dominated convergence theorem, because, using (2.9) we get $\Delta_k(y) \leq \sup_{t \in [0, 1]} F^k(t, y) \leq \int_y^{\infty} \hat{f}(1, z) dz \leq e$. Using this and taking $k \rightarrow \infty$ in (2.12) we get

$$(2.13) \quad F(t, x) = F(0, x) + \int_0^t \int_{-\infty}^{\infty} F(s, y) \rho(x - y) dy ds.$$

To finish the proof it is enough to show that the mapping $s \mapsto \int_{-\infty}^{\infty} F(s, y) \rho(x - y) dy$ is continuous, since if that is the case then we can differentiate (2.13) and deduce (FB1). This actually follows easily from (2.13):

$$\begin{aligned}
(2.14) \quad & \left| \int_{-\infty}^{\infty} F(s + h, y) \rho(x - y) dy - \int_{-\infty}^{\infty} F(s, y) \rho(x - y) dy \right| \\
&= \int_{-\infty}^{\infty} \int_s^{s+h} \int_{-\infty}^{\infty} F(r, z) \rho(y - z) \rho(x - y) dz dr dy \\
&\leq \int_s^{s+h} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \rho(y - z) \rho(x - y) dz dy dr = h. \quad \square
\end{aligned}$$

Let ν_t be the probability measure defined by $\nu_t([x, \infty)) = F(t, x)$. Since F satisfies (FB') we have that for every $b > a > \gamma(t)$,

$$\frac{d}{dt} \nu_t([a, b]) = \int_{-\infty}^{\infty} \nu_t([a - y, b - y]) \rho(y) dy$$

and thus by standard measure theory arguments we deduce that for every bounded and measurable φ with support contained in $(\gamma(t), \infty)$,

$$(2.15) \quad \frac{d}{dt} \int_{-\infty}^{\infty} \varphi(y) \nu_t(dy) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \varphi(x + y) \rho(y) \nu_t(dx) dy$$

(cf. (2.2)). Now if $A \subseteq (\gamma(t), \infty)$ has zero Lebesgue measure and the support of φ is contained in A , then the right-hand side above is 0 and we deduce that $\nu_t(A)$ is constant. Since $\nu_0(A) = \int_A f_0(x) dx = 0$, we have proved that ν_t is absolutely continuous with respect to the Lebesgue measure. We will denote its density by $f(t, \cdot)$, and we obviously have $F(t, x) = \int_x^{\infty} f(t, y) dy$.

At this point we are ready to finish the proof of Theorem 1 by showing that F satisfies the desired properties and then using the convergence of the tail distributions $F^N(t, x)$ to obtain the convergence in distribution of the process ν_t^N in $D([0, 1], \mathcal{P})$.

Proof of Theorem 1. We already know that $F^N(t, x) \rightarrow F(t, x)$ almost surely as $N \rightarrow \infty$ and that F can be written in terms of the integral of f . Now, by (2.13),

$$\int_x^{\infty} f(t, y) dy = \int_x^{\infty} f_0(y) dy + \int_0^t \int_{-\infty}^{\infty} \int_y^{\infty} f(s, z) \rho(x - y) dz dy ds$$

for $x > \gamma(t)$, so for any $h > 0$ we have

$$\begin{aligned}
(2.16) \quad & \frac{1}{h} \left[\int_{x+h}^{\infty} f(t, y) dy - \int_x^{\infty} f(t, y) dy \right] \\
&= -\frac{1}{h} \int_x^{x+h} f_0(y) dy - \int_0^t \int_{-\infty}^{\infty} \frac{1}{h} \int_{x-u}^{x+h-u} f(s, z) dz \rho(u) du ds.
\end{aligned}$$

The first integral on the right-hand side goes to $f_0(x)$ as $h \rightarrow 0$ (recall that $f_0(x)$ is continuous for $x > 0$). The second term goes to $\int_0^t \int_{-\infty}^{\infty} f(s, x - u) \rho(u) du ds$ by the

dominated convergence theorem. On the other hand, the left-hand side of (2.16) goes to $\frac{\partial F}{\partial x}(t, x)$, which equals $-f(t, x)$ for almost every $x > \gamma(t)$. We deduce that

$$(2.17) \quad f(t, x) = f_0(x) + \int_0^t \int_{-\infty}^{\infty} f(s, y) \rho(x - y) dy ds$$

for almost every x .

Now if $x_n \rightarrow x$, then by (2.17)

$$|f(t, x_n) - f(t, x)| \leq |f_0(x_n) - f_0(x)| + \int_0^t \int_{-\infty}^{\infty} f(s, y) |\rho(x_n - y) - \rho(x - y)| dy ds.$$

The first term on the right side goes to 0 as $n \rightarrow \infty$ because f_0 is continuous, while the second term goes to 0 by the continuity and boundedness of ρ and the dominated convergence theorem. We deduce that $f(t, x)$ is continuous in x and thus, in particular, (2.17) holds for every x . Thus $(f(t, x), \gamma(t))$ satisfies (FB). Moreover, since the above convergence can be achieved uniformly for t in compact intervals, it is easy to see that, if $f(t, x)$ is continuous in t for $x \neq \gamma(t)$ (as we will show next), then it is actually jointly continuous in t and x outside the curve $\{(t, \gamma(t)) : t \geq 0\}$. The fact that $f(t, x)$ is differentiable (and thus continuous) in t for $x \neq \gamma(t)$ follows easily from (2.17) by repeating the arguments in (2.14). $f(t, x) > 0$ for $x > \gamma(t)$ thanks to the facts that $f_0(x) > 0$ for $x > 0$ and that γ is non-decreasing because, according to the evolution defined by (FB), $f(t, x)$ only decreases when $x = \gamma(t)$.

The only thing left to show before turning to the proof of the convergence in distribution of ν_t^N in $D([0, 1], \mathcal{P})$ is that $(f(t, x), \gamma(t))$ is the unique solution of (FB). To do this, it is enough to show that if $(h(t, x), \sigma(t))$ is any given solution and $H(t, x) = \int_x^{\infty} h(t, y) dy$ then $H(t, x) = F(t, x)$. Indeed, if that is the case then the above arguments imply that $h(t, x)$ is jointly continuous in t and x outside the curve $\{(t, \gamma(t)) : t \geq 0\}$, and thus $h(t, x) = f(t, x)$, while (FB2) and (FB3) imply that $\sigma(t) = \gamma(t)$.

The idea of the proof will be to compare $h(t, x)$ and $f^k(t, x)$ by adapting the coupling introduced in the proof of Lemma 2.4 to the deterministic system, so we will sketch the main ideas and leave the details to the reader. We will write

$$f^k(t, x) = g^k(t, x) + d^k(t, x) + b^k(t, x),$$

which is to be interpreted in a manner analogous to (2.4). We construct these three functions inductively on each dyadic interval. We start with $g^k(0, x) = f_0(x)$ and $d^k(0, x) = b^k(0, x) = 0$ for all x . Next, for $t \in (0, \frac{1}{2^k})$ we let the three functions evolve according to the following system:

$$(2.18) \quad \begin{aligned} \frac{\partial g^k}{\partial t}(t, x) &= \int_{-\infty}^{\infty} g^k(t, y) \rho(x - y) dy && \text{for } x > \sigma(t) \\ g^k(t, x) &= 0 && \text{for } x \leq \sigma(t) \\ \frac{\partial d^k}{\partial t}(t, x) &= \mathbf{1}_{x \leq \sigma(t)} \int_{-\infty}^{\infty} g^k(t, y) \rho(x - y) dy \\ \frac{\partial b^k}{\partial t}(t, x) &= \int_{-\infty}^{\infty} [d^k(t, y) + b^k(t, y)] \rho(x - y) dy \end{aligned}$$

Then it is clear that $f^k(t, x)$ satisfies (2.3) on this interval as required. At time $\frac{1}{2^k}$ we need to shave off the extra mass in f^k . Observe that $g^k((\frac{1}{2^k})-, x) = h((\frac{1}{2^k})-, x)$ for all x , so $X_1^k \geq \sigma(\frac{1}{2^k})$. Thus all the mass to the left of $\sigma(\frac{1}{2^k})$ needs to be erased, so we put $d^k(\frac{1}{2^k}, x) = 0$ for all x . The rest of the mass to be erased from f^k will come from both g^k and b^k . This leaves us with

$$g^k(\frac{1}{2^k}, x) \leq h(\frac{1}{2^k}, x), \quad d^k(\frac{1}{2^k}, x) = 0, \quad b^k(\frac{1}{2^k}, x) \geq 0.$$

We continue the construction inductively. Assume that the above holds at time $\frac{m}{2^k}$ for some $1 \leq m < 2^k$. On the interval $(\frac{m}{2^k}, \frac{m+1}{2^k})$ we let g^k , d^k and b^k evolve according to (2.18). At time $\frac{m+1}{2^k}$ we need to shave off the extra mass in f^k , and as in the proof of Lemma 2.4 it is not hard to see that after doing that we may rebalance d^k and b^k in such a way that $d^k(\frac{m+1}{2^k}, x) = 0$ for all x and the total mass of $b^k(\frac{m+1}{2^k}, x)$ is at most the total mass of $b^k((\frac{m+1}{2^k})-, x)$. Observe that this construction preserves the inequality $g^k(t, x) \leq h(t, x)$ thanks to the observation following the proof of Proposition 2.1.

Now repeating the arguments leading to (2.5) we deduce that the total bad mass $\bar{b}_t^k = \int_{-\infty}^{\infty} b^k(t, x) dx$ satisfies $\bar{b}_t^k \leq (1 + \bar{b}_{m/2^k}^k) e^{t - \frac{m}{2^k}} - (t - \frac{m}{2^k} + 1)$, and thus

$$\bar{b}_1^k \leq \frac{e-1}{e^{2^{-k}}-1} 2^{-2k} \xrightarrow{k \rightarrow \infty} 0$$

as in (2.6). Since the sequence $(\bar{b}_{m/2^k}^k)_{m=0, \dots, 2^k}$ is increasing we deduce that

$$\sup_{m=0, \dots, 2^k} \bar{b}_{m/2^k}^k \xrightarrow{k \rightarrow \infty} 0.$$

Now let $t = \frac{l}{2^k}$ be any dyadic number in $[0, 1]$ and assume that $k \geq l$. Using the fact that $d^k(t, x) = 0$ for all $x \in \mathbb{R}$ we can write

$$(2.19) \quad 0 = \int_x^\infty (h(t, y) - f^k(t, y)) dy = \int_x^\infty (h(t, y) - g^k(t, y)) dy - \int_x^\infty b^k(t, y) dy$$

so $\int_x^\infty (h(t, y) - g^k(t, y)) dy \leq \bar{b}_t^k \rightarrow 0$. Thus, since $h(t, y) \geq g^k(t, y)$, we deduce that $\int_x^\infty g^k(t, y) dy \rightarrow \int_x^\infty h(t, y) dy$ as $k \rightarrow \infty$ uniformly in x , which in turn implies by (2.19) that

$$\lim_{k \rightarrow \infty} F^k(t, x) = \lim_{k \rightarrow \infty} \int_x^\infty f^k(t, y) dy = \int_x^\infty h(t, y) dy = H(t, x)$$

uniformly in x and for every dyadic number $t \in [0, 1]$. Since $(H(t, x), \sigma(t))$ solves (FB'), H is continuous in t and we deduce that $H(t, x) = F(t, x)$ for all $t \in [0, 1]$.

To show that the sequence of processes ν_t^N converges in distribution in $D([0, 1], \mathcal{P})$ to the deterministic process ν_t in this space defined by having its densities evolve according to (FB), it is enough to prove that this sequence is tight. In fact, if $\nu_t^{N_k}$ is any convergent subsequence, then its limit ν_t is completely defined by its tail distribution at each time t , which we know must be $F(t, \cdot)$. To show that ν_t^N is tight it is enough, by Theorem 2.1 of Roelly-Coppoletta (1986), to show that for any continuous and bounded function φ on $\mathbb{R}$ the sequence of real-valued processes $\langle \nu_t^N, \varphi \rangle = \int_{-\infty}^{\infty} \varphi(x) \nu_t^N(dx)$ is tight in $D([0, T], \mathbb{R})$. Fix one such function φ . By Aldous' criterion (which we take from Theorem 2.2.2 in Joffe and Métivier (1986) and the corollary that precedes it in page 34), we need to prove that the following two conditions hold:

(i) For every rational $t \in [0, T]$ and every $\varepsilon > 0$, there is an $L > 0$ such that

$$\sup_{N>0} \mathbb{P}(|\langle \nu_t^N, \varphi \rangle| > L) \leq \varepsilon.$$

(ii) If $\mathfrak{T}_T^N$ is the collection of stopping times with respect to the natural filtration associated to $\langle \nu_t^N, \varphi \rangle$ that are almost surely bounded by T , then for every $\varepsilon > 0$

$$\lim_{r \rightarrow 0} \limsup_{N \rightarrow \infty} \sup_{\substack{s < r \\ \tau \in \mathfrak{T}_T^N}} \mathbb{P}\left(\left|\langle \nu_{(\tau+s) \wedge T}^N, \varphi \rangle - \langle \nu_\tau^N, \varphi \rangle\right| > \varepsilon\right) = 0.$$

The first condition holds trivially in our case by taking $L > \|\varphi\|_\infty$. To get the second one fix $N > 0$, $\varepsilon > 0$, $0 < s < r$ and $\tau \in \mathfrak{T}_T^N$ and let K be the number of branchings in η_t^N on the interval $[\tau, (\tau + s) \wedge T]$. Observing that

$$\left|\langle \nu_{(\tau+s) \wedge T}^N, \varphi \rangle - \langle \nu_\tau^N, \varphi \rangle\right| \leq \frac{2\|\varphi\|_\infty}{N} K$$

and $\mathbb{E}(K) \leq Ns < Nr$, we deduce by Markov's inequality that

$$\mathbb{P}\left(\left|\langle \nu_{(\tau+s) \wedge T}^N, \varphi \rangle - \langle \nu_\tau^N, \varphi \rangle\right| > \varepsilon\right) \leq \mathbb{P}\left(\frac{2\|\varphi\|_\infty}{N} K > \varepsilon\right) \leq \frac{2\|\varphi\|_\infty \mathbb{E}(K)}{\varepsilon N} < \frac{2\|\varphi\|_\infty r}{\varepsilon}$$

and (ii) follows. $\square$

3 Proof of the results for the finite system

Now we turn to the properties of the finite system. Let us first give an explicit construction of our process. Consider an i.i.d. family $(U_i^N)_{i \geq 1}$ with uniform distribution on $\{1, \dots, N\}$ and an i.i.d. family $(R_i)_{i \geq 1}$ with distribution ρ and let $(T_i)_{i \geq 1}$ be the jump times of a Poisson process with rate 1. To construct η_t^N we proceed as follows. Assume that $\eta_0^N(1) \geq \dots \geq \eta_0^N(N)$. At time T_1 , branch $\eta_{T_1}^N(U_1^N)$ using R_1 for the displacement, erase the leftmost particle, and then relabel the particles to keep the ordering. Continue this inductively, using the variables U_i^N and R_i to determine the branching at time T_i .

This construction allows us to give a monotone coupling for two copies of the process. As in the proof of Proposition 2.2, for $\mu, \nu \in \mathcal{M}$ we will say that $\mu \preceq \nu$ whenever $\mu([x, \infty)) \leq \nu([x, \infty))$ for all $x \in \mathbb{R}$. Observe that if $\mu = \sum_{i=1}^{N_1} \delta_{x_i}$ with $x_1 \geq \dots \geq x_{N_1}$ and $\nu = \sum_{i=1}^{N_2} \delta_{y_i}$ with $y_1 \geq \dots \geq y_{N_2}$ then $\mu \preceq \nu$ if and only if $N_1 \leq N_2$ and $x_i \leq y_i$ for $i = 1, \dots, N_1$. It is easy to check that if $\eta_t^{N_1}$ and $\xi_t^{N_2}$ are two copies of our process (note that we allow them to have different total number of particles) with $\eta_0^{N_1} \preceq \xi_0^{N_2}$ (i.e., in the sense that $\sum_{i=1}^{N_1} \delta_{\eta_0^{N_1}(i)} \preceq \sum_{i=1}^{N_2} \delta_{\xi_0^{N_2}(i)}$), then if we use the same branching times and displacements and the same uniform variables for the particles in $\eta_t^{N_1}$ and the leftmost N_1 particles in $\eta_t^{N_2}$, then we have $\eta_t^{N_1} \preceq \xi_t^{N_2}$ for all $t \geq 0$.

For most of the proof of Theorem 2 it will more convenient to work with the discrete time version of our process η_n^N which is defined as follows: at each time step, choose one particle uniformly at random, and then branch that particle and remove the leftmost particle among the $N+1$. The variables U_i^N and R_i can be used to decide which particle to branch and where to send its offspring at each time step.

Proof of Theorem 2(a). We will first prove that each of the two limits exists with probability 1 and in L^1 and that the limits are non-random. We will do this for the discrete time process and leave to the reader the (easy) extension to the continuous time case. We borrow the proof from that of Proposition 1.1 in Bérard and Gouéré (2008). Since it is simple we include it for convenience. Observe that it is enough to prove the result for an arbitrary deterministic initial condition $\eta_0 \in \mathcal{X}_N$, which we fix below.

Consider the variables U_i^N , R_i and T_i introduced in the coupling above. For each $k \geq 0$ let $(\eta_{k,n}^N)_{n \geq 0}$ be a copy of our process, started at η_0 , constructed as follows: if $\eta_{k,n}^N$ is given then we let $\eta_{k,n+1}^N$ be specified by adding a particle at $\eta_{k,n}^N(U_{n+k}^N) + R_{n+k}$ and then removing the leftmost particle. It is clear then that for any $k, l \geq 0$, $\max \eta_{0,k+l}^N \leq \max \eta_{0,k}^N + \max \eta_{k,l}^N$. Moreover, for any given $d \geq 1$ the family $(\eta_{dn,d}^N)_{n \geq 1}$ is i.i.d., while the distribution of $(\eta_{k,n}^N)_{n \geq 0}$ does not depend on k . Now for $0 \leq k \leq n$ define $\xi_{k,n}^N = \eta_{k,n-k}^N$. Then $\max \xi_{0,k+l}^N \leq \max \xi_{0,k}^N + \max \xi_{k,k+l}^N$, the family $(\max \xi_{dn,d(n+1)}^N)_{n \geq 1}$ is i.i.d for any $d \geq 1$ and the distribution of $(\max \xi_{k,n+k}^N)_{n \geq 0}$ does not depend on k . It is not hard to check that $\max \xi_{k,n}^N$ satisfies the rest of the hypotheses of the subadditive ergodic theorem (see Theorem 6.6.1 in Durrett (2004)) and thus $\lim_{n \rightarrow \infty} \max \xi_{0,n}^N/n$ exists almost surely and in L^1 , and moreover the limit is non-random. Since $(\xi_{0,n}^N)_{n \geq 0}$ has the same distribution as $(\eta_n^N)_{n \geq 0}$, the same holds for $\lim_{n \rightarrow \infty} \max \eta_n^N/n$.

The above proof can be straightforwardly adapted to obtain the existence of the limit for $\min \eta_t^N/t$. To show that the two limits are equal it is enough to prove that $(\max \eta_t^N - \min \eta_t^N)/t \rightarrow 0$ in probability as $t \rightarrow \infty$. Observe that if we follow the genealogy of the particle at $\max \eta_t^N$ back in time and go back in time N generations then we will necessarily reach a particle that is not in η_t^N (at time t), and that is thus to the left of $\min \eta_t^N$. If we call X_t the position of this particle and let N_t be the number of branchings in the system up to time t , then clearly $\max \eta_t^N - X_t \leq \sum_{i=N_t-N}^{N_t} |R_i|$. Thus for any $\varepsilon > 0$ we have that

$$\begin{aligned} \mathbb{P}\left(\left|\frac{\max \eta_t^N - \min \eta_t^N}{t}\right| > \varepsilon\right) &\leq \mathbb{P}\left(|R_i| > \sqrt{t} \text{ for some } N_t - N \leq i \leq N\right) + \mathbf{1}_{\frac{N\sqrt{t}}{t} > \varepsilon} \\ &= 1 - \mathbb{P}\left(|R_1| \leq \sqrt{t}\right)^N + \mathbf{1}_{\frac{N}{\sqrt{t}} > \varepsilon} \xrightarrow{t \rightarrow \infty} 0. \end{aligned}$$

The monotone coupling introduced above allows to deduce that a_N is non-decreasing. On the other hand, in the case $N = 1$ we have that $\eta_t^1(1)$ is simply a random walk jumping at rate 1 whose jump distribution is that of $R_1 \vee 0$. Therefore $\mathbb{E}(\eta_t^1(1)) = bt$ with $b = \int_0^\infty x\rho(x) dx > 0$, and thus $a_N \geq a_1 = b > 0$ for all $N \geq 1$. $\square$

To prove parts (b) and (c) of Theorem 2 we will work with the discrete time version of the shifted process: $\Delta_n^N = (\Delta_n^N(1), \dots, \Delta_n^N(N))$ with

$$\Delta_n^N(j) = \eta_n^N(j) - \eta_n^N(N).$$

Recall that Δ_n^N is a Markov chain and its transitions are the same as those of η_n^N except that after erasing the leftmost particle the N remaining particles are shifted to the left so that the new leftmost one is at the origin.

Proposition 3.1. Δ_n^N is a positive recurrent Harris chain.

Proof. Following Athreya and Ney (1978), in order to show that Δ_n^N is Harris recurrent we need to show that there is a set $A \subseteq \mathcal{X}_N$ such that

- (i) $\mathbb{P}^\xi(\tau_A < \infty) = 1$ for all $\xi \in \mathcal{X}_N$, where $\tau_A = \inf\{n \geq 0: \Delta_n^N \in A\}$.
- (ii) There exists a probability measure q on A , a $\lambda > 0$ and a $k \in \mathbb{N}$ so that $\mathbb{P}^\xi(\Delta_k^N \in B) \geq \lambda q(B)$ for all $\xi \in A$ and all $B \subseteq A$.

To achieve this, choose some $L > 0$ so that $\delta = \rho((0, L)) > 0$ and let

$$A = \{\xi \in \mathcal{X}_N: \xi(i) - \xi(i+1) \in (0, L) \text{ for } i = 1, \dots, N-1 \text{ and } \xi(N) = 0\}.$$

Then for any initial condition $\Delta_0^N \in \mathcal{X}_N$ we can get to A in $N-1$ steps via the following path: at time 1 we choose to branch the rightmost particle (the one at $\Delta_0^N(1)$, which happens with probability N^{-1}) and send the newborn particle to a location $x_1 \in (\Delta_0^N(1), \Delta_0^N(1) + L)$ (which happens with probability at least δ). Next we branch the particle at x_1 and send the newborn particle to a location $x_2 \in (x_1, x_1 + L)$ (which happens with probability at least δ/N). If we continue this for $N-1$ steps we will end up with a configuration in A , and thus

$$(3.1) \quad \mathbb{P}^\xi(\Delta_{N-1}^N \in A) \geq \left(\frac{\delta}{N}\right)^{N-1}.$$

The bound is independent of the initial condition ξ , so by the Borel-Cantelli Lemma it follows that (i) holds. Moreover, if $B \subseteq A$ is of the form $B = \{\xi \in \mathcal{X}_N: \xi(i) - \xi(i+1) \in B_i \subseteq (0, L) \text{ for } i = 1, \dots, N-1 \text{ and } \xi(N) = 0\}$, then the preceding argument implies that

$$\mathbb{P}^\xi(\Delta_{N-1}^N \in B) \geq \frac{\rho(B_1) \cdots \rho(B_{N-1})}{N^{N-1}},$$

so by taking $\lambda = N^{-N+1}$ and q to be the normalized Lebesgue measure on the first $N-1$ coordinates of the configurations in A , we deduce that (ii) also holds.

To check that Δ_n^N is positive recurrent it is enough to check that $\sup_{\xi \in \mathcal{X}_N} \mathbb{E}^\xi(\tau_A) < \infty$. This follows from (3.1) and the strong Markov property by writing

$$\begin{aligned} \mathbb{E}^\xi(\tau_A) &= \sum_{n \geq 1} \mathbb{P}^\xi(\tau_A \geq n) \leq N-1 + \sum_{i \geq 1} \sum_{n=iN}^{(i+1)N-1} \mathbb{P}^\xi(\tau_A \geq n) \\ &\leq N-1 + (N-1) \sum_{i \geq 1} \left[1 - \left(\frac{\delta}{N}\right)^{N-1}\right]^i = (N-1) \left(\frac{N}{\delta}\right)^{N-1} < \infty \end{aligned}$$

for any $\xi \in \mathcal{X}_N$. □

Proof of Theorem 2(b). The result now follows from Proposition 3.1. The fact that Δ_n^N is positive recurrent implies that the invariant measure whose existence is assured by the Harris recurrence is finite. The absolute continuity of μ_N is a direct consequence of Theorem 2(c), which we prove below, together with the fact that if the initial condition for Δ_t^N is absolutely continuous, then so is the distribution of the process at all times. □

Proof of Theorem 2(c). Let $A \subseteq \mathcal{X}_N$ and $k = N-1$ be the objects which we found satisfy (i) and (ii) in the proof of Proposition 3.1. It is enough to prove the result along the k subsequences of the form $(\Delta_{km+j}^N)_{m \geq 0}$ with $0 \leq j < k$. Moreover, using

the Markov property at time j we see that it is enough to prove the result along the subsequence $(\Delta_{km}^N)_{m \geq 0}$, which is an aperiodic recurrent Harris chain. The result for this subsequence follows from Theorem 4.1(ii) in [Athreya and Ney \(1978\)](#) as long as we have that $\sup_{\xi} \mathbb{P}_{\xi}^{\xi}(\tau_A > t) < 1$ for some $t > 0$, where τ_A is the hitting time of A . This follows easily from the estimate in (3.1) (which is uniform in ξ). $\square$

4 Proof of Theorem 4

Recall that in this part we are assuming that ρ (and hence its tail distribution R) has exponential decay and, consequently, that the moment generating function of ρ , $\phi(\theta) = \int_{-\infty}^{\infty} e^{\theta x} \rho(x) dx$, is finite for $\theta \in (-\Theta, \Theta)$ (see (1.2) and (1.7)). Before getting started with the proof of Theorem 4 we need to prove the claim we implicitly made in (1.6).

Lemma 4.1.

$$\min_{\lambda \in (0, \Theta)} \frac{\phi(\lambda)}{\lambda} = a,$$

where a is the asymptotic speed defined in Theorem 3. Moreover, letting $\lambda^* \in (0, \Theta)$ be the number such that $\phi(\lambda^*)/\lambda^* = a$, we have that $\phi'(\lambda^*) = a$, $\phi(\lambda)/\lambda$ is strictly convex on $(0, \Theta)$ and the sign of $\phi'(\lambda) - \phi(\lambda)/\lambda$ equals that of $\lambda - \lambda^*$.

Proof. Define $c(\lambda) = \phi(\lambda)/\lambda$ for $\lambda \in (0, \lambda^*)$. A little calculus shows that $c(\lambda)$ is strictly convex:

$$c''(\lambda) = \frac{\phi''(\lambda)}{\lambda} - \frac{2\phi'(\lambda)}{\lambda^2} + \frac{2\phi(\lambda)}{\lambda^3} = \frac{1}{\lambda^3} \int_{-\infty}^{\infty} [(\lambda x - 1)^2 + 1] e^{\lambda x} \rho(x) dx > 0.$$

It is clear that $c(\lambda) \rightarrow \infty$ as $\lambda \rightarrow 0$. On the other hand, (1.3) implies $c(\lambda) \rightarrow \infty$ as $\lambda \rightarrow \Theta-$ as well. Thus the minimum of c is attained at some $\lambda^* \in (0, \Theta)$, and we have $c'(\lambda^*) = 0$, or $\phi'(\lambda^*) = \phi(\lambda^*)/\lambda^*$, which will give the second claim in the lemma once we show that $c(\lambda^*) = a$.

Now recalling the definition of a we need to show that

$$\sup_{\theta > 0} [\theta c(\lambda^*) - \phi(\theta)] = 0.$$

This is easy: using the definition of c we get

$$\sup_{\theta > 0} [\theta c(\lambda^*) - \phi(\theta)] \geq \lambda^* c(\lambda^*) - \phi(\lambda^*) = 0,$$

while for all $\theta > 0$

$$\theta c(\lambda^*) - \phi(\theta) \leq \theta c(\theta) - \phi(\theta) = 0.$$

Finally, to get the last claim in the lemma recall that $c'(\lambda^*) = 0$ and c is convex, so

$$\phi'(\lambda) - \frac{\phi(\lambda)}{\lambda} = \lambda c'(\lambda)$$

is negative for $\lambda \in (0, \lambda^*)$ and positive for $\lambda \in (\lambda^*, \Theta)$. $\square$

As we explained in Section 1.3 the proof of Theorem 4 will depend on looking for positive solutions to the equation (1.8). We will actually consider a slightly more general equation:

$$(4.1) \quad U(x) = \int_0^\infty U(y)k(x-y)dy \quad \forall x \geq 0,$$

where we look for a non-decreasing solution U , continuous except at the origin, with $U(x) = 0$ for all $x < 0$. In Spitzer's terminology, a solution U with the above properties is a P^* -solution of (4.1). When $\lim_{x \rightarrow \infty} U(x) = 1$ we call U a P -solution, and think of it as the distribution function of a non-negative random variable. Here are the two results of Spitzer that we will need.

Theorem 4.2 (Theorems 2 and 4 in Spitzer (1957)).

- (a) If $\int_{-\infty}^\infty xk(x)dx \leq 0$ then there is a unique (up to multiplicative constant) P^* -solution of (1.5).
- (b) If $\int_{-\infty}^\infty xk(x)dx < 0$ then there is a unique P -solution of (1.5) which can be obtained as the limit $U(x) = \lim_{n \rightarrow \infty} U_n(x)$ of the iterative procedure defined by $U_{n+1}(x) = \int_0^\infty U_n(y)k(x-y)dy$ starting with an arbitrary continuous U_0 corresponding to the distribution function of a non-negative random variable.
- (c) If $\int_{-\infty}^\infty xk(x)dx \geq 0$ then (1.5) has no P -solution.

Repeating the arguments we used to show that $F(t, x)$ had a density (see (2.15)) we see that if U is a P^* -solution of (4.1) then there is a non-negative function u such that $U(x) = \int_0^x u(y)dy$. Again repeating previous arguments (see the first part of the proof of Theorem 1), we deduce that u satisfies (1.8), while obviously $u(x) = 0$ for $x \leq 0$. u is continuous except possibly at the origin by the dominated convergence theorem thanks to the fact that k is continuous. Multiplying $u(x)$ by $e^{\lambda x}$ will allow us to obtain a solution for (1.5) with the desired properties.

Proof of Theorem 4. Let $c \geq a$ and take $\lambda \in (0, \lambda^*]$ such that $\phi(\lambda)/\lambda = c$ as above. The uniqueness of the solutions of (1.5) in this case follows from Theorem 4.2(a). In fact, if w_1 and w_2 are two solutions of (1.5) then $u_i(x) = e^{\lambda x}w_i(x)$ solves (1.8) for $i = 1, 2$, and thus the functions $U_i(x) = \int_0^x u_i(y)dy$ are P^* -solutions of (4.1), and they are continuous because the u_i are locally integrable. Hence $U_1(x) = AU_2(x)$ for all $x \in \mathbb{R}$ and some $A > 0$, which implies that $w_1(x) = Aw_2(x)$ for all $x \in \mathbb{R}$. Integrating this relation we get $A = 1$ and uniqueness follows.

To show existence we start by integrating by parts to obtain

$$\int_{-\infty}^\infty xk(x)dx = \frac{\lambda}{\phi(\lambda)} \int_{-\infty}^\infty xe^{\lambda x}R(x)dx = \frac{1}{\phi(\lambda)} \int_{-\infty}^\infty \left(x - \frac{1}{\lambda}\right) e^{\lambda x} \rho(x) dx.$$

Thus the sign of $\int_{-\infty}^\infty xk(x)dx$ is the same as that of $\phi'(\lambda) - \phi(\lambda)/\lambda$. This last quantity is strictly negative for $\lambda \in (0, \lambda^*)$ and vanishes for $\lambda = \lambda^*$ by Lemma 4.1.

If $c > a$, then $\lambda < \lambda^*$ and thus Theorem 4.2(b) provides us with a P -solution of (4.1) to which, by the discussion preceding this proof, we can associate a continuous (except at the origin) function u satisfying (1.8) and corresponding to the density of a non-negative random variable. Now let $A = \int_0^\infty e^{-\lambda x}u(x)dx$ which is clearly finite (actually $A < 1$). Define $w(x) = A^{-1}e^{-\lambda x}u(x)$. Then w is the density of a non-negative

random variable and it is easy to check that it satisfies (1.5a) with $c = \phi(\lambda)/\lambda$. w obviously satisfies (1.5b), (1.5c) and (1.5d) as well, and it is continuous except possibly at the origin because so is u and, by definition, $\int_0^\infty e^{\lambda x} w(x) dx = A^{-1} < \infty$. The last thing left to show in this case is that w is differentiable (except at the origin), but this follows easily from writing, for $x \geq 0$,

$$w(x) = \frac{1}{c} \int_0^\infty w(y) \int_{x-y}^\infty \rho(z) dz dy = \frac{1}{c} \int_x^\infty \int_0^\infty w(y) \rho(z-y) dy dz,$$

and using the fact that the integrand above is continuous.

To prove the second claim in (b) we may obviously assume that $\lambda' \in (\lambda, \lambda^*)$. Suppose that $\tilde{w}$ solves (1.5) and let $\tilde{W}(x) = \int_x^\infty \tilde{w}(y) dy$. It is not hard to check that $\tilde{W}$ satisfies

$$\tilde{W}(x) = \frac{1}{c} \int_{-\infty}^\infty \tilde{W}(y) R(x-y) dy \quad \forall x \geq 0.$$

Then if $A = \sup_{x \geq 0} e^{\tilde{\lambda} x} \tilde{W}(x) < \infty$ we have that

$$\begin{aligned} \tilde{W}(x) e^{\tilde{\lambda} x} &= \frac{1}{c} \int_{-\infty}^\infty \tilde{W}(y) e^{\tilde{\lambda} y} e^{\tilde{\lambda}(x-y)} R(x-y) dy \leq \frac{A}{c} \int_{-\infty}^\infty e^{\tilde{\lambda}(x-y)} R(x-y) dy \\ &= \frac{A}{c} \int_{-\infty}^\infty e^{\tilde{\lambda} u} R(u) du = \frac{A}{c} \frac{\phi(\tilde{\lambda})}{\tilde{\lambda}}. \end{aligned}$$

Taking supremum in $x \geq 0$ and recalling that $c = \phi(\lambda)/\lambda$ the above says that

$$A \leq \frac{\phi(\tilde{\lambda})}{\tilde{\lambda}'} \frac{\lambda}{\phi(\lambda)} A,$$

and since $A > 0$ this says that $\phi(\tilde{\lambda})/\tilde{\lambda} \geq \phi(\lambda)/\lambda$. But Lemma 4.1 implies exactly the opposite. This is a contradiction, and thus $A = \infty$.

The case $c = a$ is similar so we will skip some details. Now we have $\lambda = \lambda^*$ and thus $\int_{-\infty}^\infty x k(x) dx = 0$. Theorem 4.2(a) provides us now with a P^* -solution of (1.5) to which corresponds a function u which is the density of a measure supported on $[0, \infty)$ and which satisfies (1.8). Let $A = \int_0^\infty e^{-\lambda^* x} u(x) dx$. We need this quantity to be finite, so that $w(x) = A^{-1} e^{-\lambda^* x} u(x)$ be a (continuous) probability density. This follows from Theorem 6.2 of Engibaryan (1996), which assures that $U(x) \leq Cx$ for some $C > 0$, and integration by parts:

$$\begin{aligned} A &= \int_0^\infty e^{-\lambda^* x} u(x) dx = \lim_{x \rightarrow \infty} e^{-\lambda^* x} U(x) - U(0) + \lambda^* \int_0^\infty e^{-\lambda^* x} U(x) dx \\ &\leq C \lambda^* \int_0^\infty e^{-\lambda^* x} x dx < \infty. \end{aligned}$$

It is easy again to verify that w satisfies (1.5), and its differentiability follows from the same reasons as above. Clearly $\int_0^x e^{\lambda^* y} w(y) dy = A^{-1} \int_0^x u(y) dy = O(x)$, so w is a solution with the desired properties.

We are only left showing (d), that is, that there are no solutions of (1.5) when $c < a$. We start by observing that if w were a solution then the above arguments would imply that w is differentiable on the positive axis, and thus if we set $f_0(x) = w(x)$ in (FB) we

get $\gamma(t) = ct$. Therefore, to show the non-existence of solutions for $c < a$ it is enough to show that, given any $\varepsilon > 0$ and any f_0 supported on $[0, \infty)$, there is a $T > 0$ such that the solution $(f(t, x), \gamma(t))$ of (FB) satisfies

$$(4.2) \quad \gamma(T) > (a - \varepsilon)T.$$

To achieve this consider the following construction. Take $M \geq 1$ and let χ_0 be such that $\int_{\chi_0}^{\infty} f_0(x) dx = 1/M$. Now recall the definition of the process $\hat{\nu}_t$ in the proof of Proposition 2.1, which corresponded to the deterministic measure valued limit of $\hat{\nu}_t^N$, and observe that we can run this process started with any initial measure (not necessarily an absolutely continuous one). Moreover, (2.2) still holds in this case by Theorem 5.3 of Fournier and Méléard (2004). Letting $\hat{\nu}_0 = M^{-1}\delta_{\chi_0}$ (a point mass at χ_0 of size $1/M$), run the process $\hat{\nu}_t$ up to time $t = \log(M)$ and define χ_1 to be such that $\hat{\nu}_{\log(M)}([\chi_1, \infty)) = 1/M$.

We need to prove two things, which together will imply (4.2). The first one is that $\gamma(\log(M)) \geq \chi_1$. To see that this is indeed the case, consider a copy of the process ν_t started at ν_0 (which is the probability measure corresponding to f_0) and a copy of $\hat{\nu}_t$ started at $M^{-1}\delta_{\chi_0}$. It is clear that $\hat{\nu}_0 \preceq \nu_0$, in the sense of the ordering introduced in Section 3, and since the mass of $\hat{\nu}_t$ is bounded by 1 for $t \in [0, \log(M)]$, the evolution of these processes preserves this inequality up to time $\log(M)$. This implies that $\chi_1 \leq \gamma(\log(M))$. The above arguments can be made precise using some of the ideas introduced in the proof of Theorem 1, but we leave the details to the reader.

Next we need to prove that

$$(4.3) \quad \chi_1 > (a - \varepsilon) \log(M)$$

as long as M is large enough. This together with the bound of the preceding paragraph imply (4.2) by taking $T = \log(M)$. To show (4.3) we may assume, by the translation invariance of $\hat{\nu}_t$, that $\chi_0 = 0$. Observe that $\hat{\nu}_t$, started with $\hat{\nu}_0 = \delta_0$, corresponds to the mean measure of the branching random walk ξ_t^1 (started with just one particle located at the origin). This can be made precise by roughly writing down the formula for the generator of ξ_t^1 and applying it to functions of the form $\xi \mapsto \langle \xi, \varphi \rangle = \sum_{i=1}^{N(\xi)} \varphi(\xi(i))$, where $N(\xi)$ is the number of particles in the branching random walk configuration ξ , to deduce that after taking expectations the resulting equation is the same as (2.2). We leave the details to the reader, and instead only state that the above implies that

$$(4.4) \quad \hat{\nu}_t([ct, \infty)) = \mathbb{E}(\xi_t^1([ct, \infty)))$$

for every c , where $\xi_t^1([ct, \infty))$ denotes the number of particles in the branching random walk to the right of ct at time t . On the other hand it is well known that

$$\mathbb{E}(\xi_t^1([ct, \infty))) = e^t \mathbb{P}(S_t \geq ct)$$

where S_t is defined as in (1.4), and thus (1.4) implies that

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log(\mathbb{E}(\xi_t^1([ct, \infty)))) = \Lambda(c) + 1.$$

Since Λ is strictly decreasing on $[0, \infty)$ and $\Lambda(a) = -1$ we deduce that

$$a = \sup \left\{ c > 0 : \lim_{t \rightarrow \infty} \frac{1}{t} \log(\mathbb{E}(\xi_t^1([ct, \infty)))) > 0 \right\}.$$

This together with (4.4) implies that

$$(4.5) \quad \widehat{\nu}_t([(a - \varepsilon)t, \infty)) > 1$$

for large enough t . On the other hand, if we start $\widehat{\nu}_t$ at $M^{-1}\delta_0$, then $\chi_1 > (a - \varepsilon)\log(M)$ if and only if $\widehat{\nu}_{\log(M)}([(a - \varepsilon)\log(M), \infty)) > 1/M$. This is equivalent to starting $\widehat{\nu}_t$ at δ_0 and asking for $\widehat{\nu}_{\log(M)}([(a - \varepsilon)\log(M), \infty)) > 1$, which holds for large enough M by (4.5). We have obtained (4.3) and thus the proof is complete. $\square$

References

- Athreya, K. B. and Ney, P. (1978). A new approach to the limit theory of recurrent Markov chains. *Trans. Amer. Math. Soc.*, **245**, 493–501.
- Bérard, J. and Gouéré, J.-B. (2008). Brunet-derrida behavior of branching-selection particle systems on the line. arXiv:0811.2782.
- Biggins, J. D. (1977). Chernoff’s theorem in the branching random walk. *J. Appl. Probability*, **14**(3), 630–636.
- Brunet, E. and Derrida, B. (1997). Shift in the velocity of a front due to a cutoff. *Phys. Rev. E* (3), **56**(3, part A), 2597–2604.
- Chayes, L. and Swindle, G. (1996). Hydrodynamic limits for one-dimensional particle systems with moving boundaries. *Ann. Probab.*, **24**(2), 559–598.
- Durrett, R. (2004). *Probability: theory and examples*. Duxbury Press, Belmont, CA, third edition.
- Durrett, R. and Mayberry, J. (2008). Evolution in predator-prey systems. Preprint.
- Engibaryan, N. B. (1996). Convolution equations containing singular probability distributions. *Izv. Ross. Akad. Nauk Ser. Mat.*, **60**(2), 21–48.
- Feller, W. (1971). *An introduction to probability theory and its applications. Vol. II*. Second edition. John Wiley & Sons Inc., New York.
- Fournier, N. and Méléard, S. (2004). A microscopic probabilistic description of a locally regulated population and macroscopic approximations. *Ann. Appl. Probab.*, **14**(4), 1880–1919.
- Gravner, J. and Quastel, J. (2000). Internal DLA and the Stefan problem. *Ann. Probab.*, **28**(4), 1528–1562.
- Joffe, A. and Métivier, M. (1986). Weak convergence of sequences of semimartingales with applications to multitype branching processes. *Adv. in Appl. Probab.*, **18**(1), 20–65.
- Kreĭn, M. G. (1962). Integral equations on a half-line with kernel depending upon the difference of the arguments. *Transl. Amer. Math. Soc. (2)*, **22**, 163–288. Original (Russian): *Uspekhi Mat. Nauk.*, **13**(5) (1958), 3–120.

- Meirmanov, A. M. (1992). *The Stefan problem*, volume 3 of *de Gruyter Expositions in Mathematics*. Walter de Gruyter & Co., Berlin. Translated from the Russian by Marek Niezgódka and Anna Crowley, With an appendix by the author and I. G. Götz.
- Paley, R. E. A. C. and Wiener, N. (1987). *Fourier transforms in the complex domain*, volume 19 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI. Reprint of the 1934 original.
- Roelly-Coppoletta, S. (1986). A criterion of convergence of measure-valued processes: application to measure branching processes. *Stochastics*, **17**(1-2), 43–65.
- Spitzer, F. (1957). The Wiener-Hopf equation whose kernel is a probability density. *Duke Math. J.*, **24**, 327–343.
- Wiener, N. and Hopf, E. (1931). Über eine klasse singulärer integralgleichungen. *Sitzungber. Akad. Wiss. Berlin*, page 696706.